



# Cheetah<sup>®</sup> extra >>>

**CONTAINS** An oil in water emulsion containing 69 g/L (7.14% w/w) fenoxaprop-p-ethyl.

**CONTROLS** For use only as an agricultural herbicide for the control of wild-oats and grass-weeds in winter and spring wheat.

PCS 04168

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## RATE OF USE

CROP	MAXIMUM INDIVIDUAL DOSE	MAXIMUM TOTAL DOSE	LATEST TIME OF APPLICATION
Winter wheat, spring wheat	1.0 L product per hectare	1.0 L product per hectare	From emergence of the crop up to and including flag leaf fully emerged (GS39) but before flag leaf sheath extending stage (GS41)

## RESTRICTIONS

**DO NOT** apply Cheetah Extra to barley, durum wheat, undersown crops or those to be undersown.

**DO NOT** apply Cheetah Extra to wheat beyond the flag leaf ligule visible stage (GS 39) of the crop.

**DO NOT** roll or harrow within one week of spraying.

**DO NOT** spray crops under stress, suffering drought, waterlogged, grazed, lacking nutrients or if soil is compacted.

Broadcast crops should be sprayed post-emergence after the plants have a well-established root system and all other conditions on the label and leaflet are met.

Cheetah Extra can be sprayed in frosty weather provided the crop has hardened off.

Cheetah Extra is rainfast one hour after spraying.

**DO NOT** spray wet foliage liable to run off or leaves covered in ice.

Avoid spraying immediately before or after a sudden or extended drop in temperature or a period of warm days/cold nights.

Dry conditions resulting in moisture stress may reduce effectiveness. Such conditions will tend to occur with late applications to spring crops.

## WEEDS CONTROLLED

### **Wild-oats, rough meadow-grass and awned canary-grass.**

Strains of some annual grasses (e.g. black-grass, wild-oats and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

Cheetah Extra will control black-grass and wild-oats at the 2 fully expanded leaf stage (GS 12) up to and including the flag leaf ligule just visible (GS 39).

Cheetah Extra will control awned canary-grass at the 2 fully expanded leaf stage (GS 12) up to the end of tillering but before first node detectable (GS 31).

Cheetah Extra will control rough meadow-grass at the 2 fully expanded leaf stage (GS12) up to the end of tillering but before first node detectable (GS31).

Cheetah Extra controls emerged weeds on all soils and tilths and is unaffected by high organic matter or high Kd factor (i.e. soils with a high adsorptive capacity).

The rates of use of Cheetah Extra depend on weed type, size and time of application.

**Wild-oats****0.68 L/ha**

Apply from when the weeds have at least 2 fully expanded leaves (GS 12) up to the early tillering stage (GS 23).

**Late-drilled crops**

Susceptible weeds emerging in the spring in late-drilled crops (i.e. drilled during November and December) will be controlled with a spring application of Cheetah Extra at 0.68 L/ha provided they have at least two fully expanded leaves (GS 12) and are no larger than the early tillering stage (GS 23). For larger weeds, use the appropriate higher rate.

**0.8 L/ha**

Apply for the control of weeds at the 2 fully expanded leaf stage (GS 12) up to the end of tillering stage but before first node detectable (GS 31). For weeds beyond the end of tillering, always use the 1.0 L/ha rate.

**1.0 L/ha**

Apply for the control of weeds at the 2 fully expanded leaf stage (GS 12) up to and including the flag leaf ligule just visible stage (GS 39).

**Awned canary-grass and rough meadow-grass****0.68 L/ha**

Apply from when the weeds have at least 2 fully expanded leaves (GS 12) up to the early tillering stage (GS 23), normally until the end of December.

**0.8 L/ha**

Apply from when the weeds have at least 2 fully expanded leaves (GS 12) up to the end of tillering stage but before first node detectable (GS 31).

**Maximum total dose:** 1.0 L/ha

**CROP SPECIFIC INFORMATION**

For use on Winter wheat, Spring wheat.

Apply Cheetah Extra from emergence of the crop up to and including flag leaf fully emerged (GS 39) but before flag leaf sheath extending stage (GS 41).

Cheetah Extra is a foliar-acting herbicide. For optimal results, use application techniques which ensure good weed coverage and crop penetration, using flat fan nozzles.

**Conventional Volume**

**Volume:** 200-300 litres of water per hectare

Always use the higher volume when weeds or crop are dense to avoid shading of the weed.

**Pressure:** 2.5-3.0 bars

**Spray Quality:** FINE TO MEDIUM (BCPC category)

## **MIXING**

### **SHAKE WELL BEFORE USE**

Add the recommended quantity of Cheetah Extra to the spray tank half-filled with the required quantity of clean water. Add the remainder of the water with the sprayer agitation system in operation. When emptying containers, wash out and add washings to the sprayer. Agitate before and during spraying. **DO NOT** leave the sprayer standing with chemical in it. After spraying, thoroughly wash out the spray tank.

## **FOLLOWING CROPS AND SEQUENCES**

Cheetah Extra can be used as part of a spray programme with any other spray chemical provided an interval of at least 7 days elapses between applications.

## **COMPATIBILITY**

Cheetah Extra may be applied as a tank-mix with a range of products. Contact Bayer CropScience for compatibility information on specific tank-mixes. Full manufacturer's instructions must be followed for each tank-mix component.

Continuous agitation must be maintained.

**DO NOT** mix Cheetah Extra with herbicides containing phenoxy or sulfonylurea containing products other than Eagle.

## SAFETY PRECAUTIONS

### Operator Protection

**WEAR SUITABLE PROTECTIVE CLOTHING GLOVES AND EYE/FACE PROTECTION** when handling the concentrate.

**WASH HANDS** and exposed skin before meals and after use

### Environmental Protection

**DO NOT CONTAMINATE PONDS, WATERWAYS OR DITCHES** with chemical or used container.

### Storage and Disposal

**DO NOT REUSE CONTAINER** for any other purpose and dispose of safely.

Store in a safe dry place designated as an agrochemical store.


**PROTECT FROM FROST.**

**READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.**

## RISK AND SAFETY INFORMATION

### CHEETAH EXTRA

Contains 69 g/L (7.14% w/w) fenoxaprop-P-ethyl.



**IRRITANT**



**DANGEROUS  
FOR THE  
ENVIRONMENT**

**MAY CAUSE SENSITISATION BY SKIN CONTACT.  
TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS  
IN THE AQUATIC ENVIRONMENT.  
REPEATED EXPOSURE MAY CAUSE SKIN DRYNESS OR CRACKING.**

Keep out of reach of children.

Keep away from food, drink and animal feedingstuffs.

Avoid contact with skin.

Wear suitable gloves.

This material and its container must be disposed of in a safe way.

Use appropriate containment to avoid environmental contamination.

To avoid risks to man and the environment, comply with the instructions for use.

### PACK SIZE

5 L