



A ketoenol insecticide for control of aphids and whitefly in Brussels sprouts, broccoli/calabrese, cabbage, cauliflower, collards and kale and aphids in lettuce.

An oil dispersion formulation containing 150 g/L (15.94% w/w) spirotetramat.

Approval Holder:
Bayer CropScience Limited
230 Cambridge Science Park
Milton Road, Cambridge CB4 0WB

Telephone
from the UK: 01223 226500
from the ROI: +44 1223 226500

For 24 hour emergency information contact
Bayer CropScience Limited
from the UK: 0800 220876
from the ROI: 1 800 409 399

Risk and Safety Information

MOVENTO

Contains 150 g/L (15.94%) spirotetramat



IRRITANT



DANGEROUS FOR THE ENVIRONMENT

May cause sensitisation by skin contact

Toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment

Keep out of reach of children
Keep away from food, drink and animal feedingstuffs
Avoid contact with skin
Wear suitable protective gloves
Use appropriate containment to avoid environmental contamination.
To avoid risks to man and the environment, comply with the instructions for use.
This material and its container should be disposed of in a safe way.

MAPP 14446 PCS 04101

SAFETY PRECAUTIONS – All UK Only

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Environmental Protection

Do not contaminate water with product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards or roads.

RISK TO NON-TARGET INSECTS OR OTHER ARTHROPODS. See directions for use.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION:

FOR USE ONLY AS A HORTICULTURAL INSECTICIDE

Crops:	Brussels sprouts, broccoli/calabrese, cabbage, cauliflower, collards, kale and lettuce
Maximum individual dose:	0.5 L/ha
Maximum number of treatments:	2
Latest time of application:	Brussels sprouts, broccoli/calabrese, cabbage, cauliflower, collards and kale: 3 days before harvest Lettuce: 7 days before harvest

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

GB79093748x rA3

DIRECTIONS FOR USE

IMPORTANT: This leaflet is approved as part of the label. All instructions on this leaflet and on the label should be read carefully in order to obtain successful results from the use of this product.

RESTRICTIONS

Shake well before use.

Avoid spraying within 5m of the field boundary to reduce effects on non-target insects or other arthropods.

PESTS CONTROLLED

Movento is a ketoenol systemic insecticide which works by inhibition of lipid biosynthesis to control the sucking pests listed below:

- Mealy cabbage aphid, *Brevicoryne brassicae*
- Peach potato aphid, *Myzus persicae*
- Blackcurrant-lettuce aphid, *Nasonovia ribis-nigri*
- Lettuce root aphid, *Pemphigus bursarius*
- Brassica whitefly, *Aleyrodes proletella*

Due to the mode of action, rapid knockdown of pests should not be expected. Obvious control usually occurs after 3-7 days and is dependent upon pest stage, with youngest larvae being most susceptible and adults least susceptible. For optimum control, application should be made as soon as pests appear in the crop, so that population build up is prevented through targeting the very young larval stages as they are produced.

Through its two-way systemicity, the active ingredient spirotetramat is translocated to growing points and is able to reach hidden aphids in heart leaves and on roots. However, it is important to note that there is little movement from new leaves to older leaves, so good spray cover on older leaves will be particularly important.

CROP SPECIFIC INFORMATION

Application

Apply in a water volume of 300 - 600 Litres of water per hectare according to crop density.

Apply as a MEDIUM quality spray (BCPC).

All varieties of Brussels sprouts, broccoli/calabrese, cabbage, cauliflower, collards and kale, and outdoor and protected lettuce may be treated.

Apply Movento at a rate of 0.5 L/ha as aphids or whitefly start to build up in the crop.

Movento may be used up to a maximum of two applications.

PESTICIDE RESISTANCE MANAGEMENT STRATEGY

Total reliance on one pesticide will hasten the development of resistance; pesticides of different chemical types or alternative control measures should be included in a planned programme. In a spray programme Movento should be used with other insecticides of a different mode of action, either in alternation or as a 2-spray block within the programme e.g. following a neonicotinoid seed or soil treatment.

Movento should always be applied at the full recommended rate of use and in sufficient water volume to achieve the required spray penetration into the crop and uniform coverage necessary for optimal pest control.

MIXING AND SPRAYING

Shake well before use. Add the required quantity of Movento to the half-filled spray tank with the agitation system in operation, and fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing. Wash equipment thoroughly, immediately after use, by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Processing Information:

If the crop is intended for processing consult the processor before the use of Movento

© Bayer CropScience 2010