

Invert pack repeatedly and shake before use!



Betanal[®] maxxPro[®]

5 L e

H

Herbicide

Betanal maxxPro is a broad-spectrum post-emergence contact/residual herbicide which controls a wide range of annual broad-leaved weeds and annual meadow-grass in sugar beet, fodder beet and mangels.

An oil dispersion containing
47 g/L (4.48 % w/w) desmedipham,
75 g/L (7.23 % w/w) ethofumesate,
27 g/L (2.56 %w/w) lenacil and
60 g/L (5.73 % w/w) phenmedipham.

For Professional use only

Authorisation holder:
Bayer CropScience Ltd.
230 Cambridge Science
Park
Milton Road
Cambridge
CB4 0WB
United Kingdom

Marketing company:
Bayer CropScience Ltd.
Bayer Ltd
The Atrium
Blackthorn Road
Sandyford
Dublin 18

Freephone: 1800 818534
For 24 hour emergency information contact
Bayer CropScience Limited Telephone: 00800 1020 3333

Safety information

BETANAL MAXXPRO

Contains 47 g/L (4.48 % w/w) desmedipham,
75 g/L (7.23 % w/w) ethofumesate,
27 g/L (2.56 %w/w) lenacil and
60 g/L (5.73 % w/w) phenmedipham.



Danger

Causes serious eye damage.

May cause an allergic skin reaction.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 04266

IE85785214b rA1j

PROTECT FROM FROST
SHAKE WELL BEFORE USE



SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

AVOID ALL CONTACT WITH SKIN.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WHEN USING DO NOT EAT DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.



betanamaxprods

To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.ie/sds/betanamaxpro.pdf

or alternatively contact your supplier

IE85828649b rA1j

Bayer

DIRECTIONS FOR USE

For Professional use only via tractor mounted sprayer.

INTRODUCTION

Betanal maxxPro is a selective weedkiller for the post-emergence control of many annual weeds in beet crops.

Betanal maxxPro is recommended for low volume overall spraying in a planned spray programme and can follow an Approved pre-emergence herbicide and / or Approved post-emergence herbicide treatment(s) used at recommended low doses for programmes.

Betanal maxxPro acts mainly by contact action but will, in a planned programme and in moist conditions, provide residual control of some later germinating weeds.

Crop:	Sugar beet, fodder beet and mangels
Maximum single dose (per crop):	1.5 L/ha
Maximum number of applications at max. single dose:	-
Max. total dose (per crop):	4.5 L/ha
Latest time of application:	When crop leaves meet between the rows
Other Specific Restrictions:	The maximum total dose must not exceed 1.0 kg ethofumesate per hectare in any three year period.

WEEDS CONTROLLED

The recommended timing of application is fully expanded cotyledon stage of the largest weed present and this should be followed wherever possible. The weed growth stage given below is for an occasional larger weed which will usually be controlled from a full programme of Betanal maxxPro sprays.

Weeds killed up to 6 expanded true leaves

Common field-speedwell

Fat-hen

Wild Radish, (Runch)

Weeds killed up to 4 expanded true leaves

Annual meadow-grass

Black-bindweed

Common fumitory

Field pansy

Field penny-cress

Ivy-leaved speedwell

Weeds killed up to 2 expanded true leaves

Black nightshade
Corn marigold
Groundsel
Henbit dead-nettle
Knotgrass
Pale persicaria
Red dead-nettle
Redshank
Scarlet pimpernel
Small nettle

Weeds killed at cotyledon stage

Bugloss
Charlock
Common orache
Volunteer oilseed rape
Crane's-bill (Qualified minor use recommendation)
Corn spurrey (Qualified minor use recommendation)

Other weeds controlled

Cleavers—killed up to 5 cm across or high.
Common chickweed—killed up to 10 cm across or high.
Mayweed—severe check or kill up to 2.5 cm across or high.

CROP INFORMATION

All commercially grown varieties of sugar beet, fodder beet and mangels may be treated with Betanal maxxPro.

Soil

Betanal maxxPro may be used on all soil types. On soils with more than 5% organic matter content residual activity may be reduced.

Factors Affecting Crop Tolerance

When used as recommended, Betanal maxxPro exhibits good safety to beet. Several factors may affect crop tolerance and individually may appear to have little effect, but may be important when two or more occur at the same time. (see COMPANY ADVISORY INFORMATION)

Drift

Avoid drift to areas outside those being sprayed, having due regard to the prevailing weather conditions and spray quality being used. The risk of drift is greater when a FINE quality spray, as recommended for this product, is used.

Rates of use

A programme of Betanal maxxPro low volume treatments should be applied as directed below.

1st spray

Crop and weeds at cotyledon stage

1.25 L/ha

2nd and subsequent sprays

Crop at or beyond two true leaf stage

1.5 L/ha

Timing

Crop:

The first Betanal maxxPro application may be made when the majority of crop plants have reached the fully expanded cotyledon stage.

Weeds:

The first flush of weeds should preferably not be larger than the fully expanded cotyledon stage at the time of spraying and must not be larger than the sizes stated under WEEDS CONTROLLED.

Where a pre-emergence treatment has been applied as a band spray, it will be necessary to time the first post-emergence Betanal maxxPro treatment according to the size of weeds present between the rows where there is no pre-emergence effect.

A single application of Betanal maxxPro is seldom sufficient to achieve season long weed control.

Susceptible weeds may not all be killed by the first treatment. They will usually be checked and then controlled by later applications. Further Betanal maxxPro sprays should be applied as each new flush of weeds reaches the cotyledon stage.

Successful results from a programme are achieved by applying each spray when the previous one is still showing an effect on the weeds, but a minimum of 5 days must elapse between treatments.

Volume

80 to 200 litres water per hectare.

Spray operations and Spray Quality

IMPORTANT—SHAKE THE CAN WELL BEFORE USE

1. Wash out the sprayer, spraybars and nozzles to ensure that no trace remains of previous products. This is most important after hormone and sulfonyl urea weedkillers. Contaminated hoses should be replaced by new hoses.
NB All hose connections must be secured with hose clips.
2. Check that nozzle tips are clean, undamaged, of the correct type to give a FINE quality spray, and are all of the same size, giving equal spray outputs and distribution.

3. Half fill the spray tank with clean water and start agitating. Check the agitation is operating correctly. Add the required quantity of Betanal maxxPro, mixing well by agitation, and complete filling to the final volume. Mix only sufficient chemical for immediate use and continue agitation until the tank load is used.
(See COMPATIBILITY AND SEQUENCES for any special instructions for preparing tank mixes.)
4. Ensure that the spray nozzles are set at the correct height to give even spray cover. Commence spraying, being careful to maintain correct pressure and tractor speed. Avoid overdosing or drift onto neighbouring crops.
5. After each day's work, drain the sprayer, wash thoroughly with water and liquid detergent and spray out completely. Ensure that all liquid is removed from the sprayer tank, pump and hoses. Remove nozzles, open tank and drain pump to allow free access of air to all parts of the system.

Betanal maxxPro should be applied overall as a FINE quality spray (for details see "Boom Sprayers Handbook" published by BCPC).

Even cover with a FINE quality spray is essential for good results. Take care to avoid overlapping.

COMPATIBILITY AND SEQUENCES

Compatibility

Betanal maxxPro is compatible with:

Goltix 70 SC (see directions below)

Betanal maxxPro plus Goltix 70 SC

For improved residual activity Goltix 70 SC may be added to Betanal maxxPro applications in a post-emergence programme.

Dose	Betanal maxxPro	1.25 L/ha
plus	Goltix 70 SC	1 kg/ha

Goltix 70 SC should be thoroughly dispersed in the spray tank one third full of water before adding Betanal maxxPro and filling to the mark. Continue agitation whilst spraying.

Timing As for Betanal maxxPro alone

These mixtures may be used up to three times in the crop (Do not exceed maximum total dose allowed for partner products).

Sequences

Programmes with pre-emergence treatments:-

Betanal maxxPro may be used as part of a planned programme after the use of approved pre-emergence treatments at recommended doses for programmes.

Programmes with post-emergence treatments for broad-leaved weeds:-

See crop tolerance section below.

Sequences with grass weedkillers:-

Betanal maxxPro may be used in sequence with grass weedkillers. Refer to graminicide product literature for recommended intervals.

SUCCEEDING CROPS

Beet crops may be sown at any time after the use of Betanal maxxPro. Any other crop may be sown 3 months after using Betanal maxxPro. Ploughing (mould board) to a minimum depth of 15 cm should precede preparation of a new seed bed.

COMPANY ADVISORY INFORMATION

Crop Tolerance

Several factors may affect the crop tolerance and performance of Betanal maxxPro especially when used in mixtures. Some transient crop effects may occur soon after application but when the crop is under stress, application of Betanal maxxPro can lead to crop check from which it may not fully recover. In extreme cases some loss of crop may occur.

Application of Betanal maxxPro or Betanal maxxPro mixtures is not recommended when the crop is suffering from stress. Stress can be caused by a number of circumstances, the most important of which are as follows.

Use after other Herbicides

Most beet crops are treated with more than one herbicide. In these circumstances it is not always possible to predict the interactions which may occur, and the effects of stresses in such conditions may be more severe.

Manganese Deficiency

If a crop is suffering from manganese deficiency it may be checked by Betanal maxxPro. To avoid crop check, manganese should ideally be applied to the crop first.

Lime Deficiency

Crops suffering from lime deficiency may be checked when sprayed with Betanal maxxPro. Growers should ensure that the lime status of the soil is satisfactory before drilling.

High Temperatures

When the temperature is, or is likely to be, above 21°C (70°F) on the day of spraying, application of Betanal maxxPro should be made after 5 pm otherwise crop check may occur.

If crops are subjected to substantial day to night temperature changes shortly before or after spraying, a check may occur from which the crop may not fully recover.

Frost

Crops sprayed with Betanal maxxPro and subjected to frost within seven days may be checked such that they may not fully recover.

High Light Intensity

Where the crop is already under stress, and Betanal maxxPro is sprayed in conditions of high light intensity, the crop may suffer a check from which it may not fully recover.

Wind

Strong winds can damage crops by physical buffeting and contingent damage can arise from blown soil particles. When wind damage has occurred, delay application of Betanal maxxPro until the crop has recovered.

Insect Attack

If the crop is damaged by insects, eg mangel fly, pygmy mangel beetle or flea beetle, Betanal maxxPro may cause a check from which the crop may not fully recover.

Storage

Store well away from seeds, fertilisers and animal feedingstuffs, out of direct sunlight in a safe, dry place designated as an agrochemical store.

RESISTANCE

Certain weeds may develop resistance to Bayer products. Since such circumstances are beyond our control, Bayer CropScience will be under no liability for any resulting loss or damage whatsoever.

© Betanal and maxxPro are registered Trade Marks of Bayer.

All other brands listed are Trade Marks of other Manufacturers and proprietary rights may exist.

© Bayer CropScience Limited 2018



SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

AVOID ALL CONTACT WITH SKIN.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WHEN USING DO NOT EAT DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE.



betanalmxxprodsd

To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.ie/sds/betanalmxxpro.pdf

or alternatively contact your supplier

IE85828649b rA1j

Bayer