



# CELLO<sup>®</sup>

5 L e

F

## Fungicide

For use only as an agricultural fungicide.

A fungicide for the control of stem-base, foliar and ear diseases in winter and spring wheat (also reduction of the mycotoxin deoxynivalenol), winter rye, triticale, winter and spring barley and oats.

An emulsifiable concentrate formulation containing  
100 g/L (10.3% w/w) prothioconazole,  
100 g/L (10.5% w/w) tebuconazole and  
250 g/L (26.3%w/w) spiroxamine

For Professional use only

Authorisation holder:  
Bayer CropScience Ltd.  
230 Cambridge Science Park  
Milton Road  
Cambridge  
CB4 0WB  
United Kingdom

Marketing company:  
Bayer CropScience Ltd.  
Bayer Ltd  
The Atrium  
Blackthorn Road  
Sandyford  
Dublin 18

Freephone: 1800 818534

For 24 hour emergency information contact  
Bayer CropScience Limited Telephone: 00800 1020 3333



### Safety information

#### CELLO

Contains 100 g/L (10.3% w/w) prothioconazole, 100 g/L (10.5% w/w) tebuconazole and 250 g/L (26.3% w/w) spiroxamine and N,N-Dimethyl decanamide.



#### Warning

**Causes skin irritation. May cause an allergic skin reaction. May cause respiratory irritation. Causes serious eye irritation. Harmful if inhaled. Suspected of damaging the unborn child. May cause damage to organs (Eyes) through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/eye protection/face protection. IF ON SKIN: wash with plenty of water/soap. IF INHALED: remove person to fresh air and keep comfortable for breathing. If skin irritation or rash occurs: Get medical attention/advice. Call a POISON CENTER / doctor/physician if you feel unwell. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS No. 05066



## SAFETY PRECAUTIONS

### Operator Protection

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.

Wear suitable protective clothing (coveralls) when applying the product.

Wear suitable protective clothing (coveralls) and suitable protective gloves when in contact with contaminated surfaces.

When using do not eat, drink or smoke.

Wash any contamination from eyes immediately.

Wash hands and exposed skin before meals and after work.

If you feel unwell, seek medical advice (show the label where possible).

### Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m\* to surface water bodies.

\*To reduce this buffer zone please refer to PRCD Guidance – STRIPE (Surface water Tool for Reducing the Impact of Pesticides in the Environment)

Aim spray away from water

### Storage and Disposal

Keep in original container tightly closed in a safe place.

Keep out of reach of children.

Keep away from food, drink and animal feedingsuffs.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

PROTECT FROM FROST



cellosdsie

To access the **Safety Data Sheet** for this product scan the code or use the link below:

[www.bayercropscience.ie/sds/cello.pdf](http://www.bayercropscience.ie/sds/cello.pdf)

or alternatively contact your supplier

IE86259605a rA4

Bayer

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Cello is a mixture of triazolinthione, triazole and spiroketoalmine fungicides recommended for control of a wide range of diseases on winter and spring barley, winter and spring wheat, triticale and winter rye.

## RATE OF USE

Crop	Maximum individual dose	Maximum number of applications	Latest time of application
Winter and spring wheat and winter rye	1.25 litres product per hectare	2 per season	At any stage before caryopsis watery ripe (GS 71)
Winter and spring barley, triticale and oats	1.25 litres product per hectare	2 per season	At any stage before beginning of flowering (GS 61)

Method of application: Tractor mounted sprayer

A spray pressure of 2-3 bar is recommended. Apply Cello in 200–400 litres per hectare water.

The maximum water volume is 400 litres of spray solution/ha.

Apply as a medium spray quality.

## DISEASES CONTROLLED

Disease	Wheat	Barley	Oats	Triticale	Rye
Eyespot ( <i>Oculimacula spp</i> )	R	R	R	R	R
Septoria Leaf Blotch ( <i>Mycosphaerella graminicola</i> )	MC			MC	MC
Glume Blotch ( <i>Stagonospora nodorum</i> )	MC			MC	MC
Powdery Mildew ( <i>Blumeria graminis</i> )	C	C	C	C	C
Yellow Rust	C	C		C	
Brown Rust	C	C		C	C

Disease	Wheat	Barley	Oats	Triticale	Rye
Crown Rust			C		
Tan Spot	MC				
Net blotch		C			
<i>Fusarium</i> Ear blight	R	MC		R	R
Sooty Moulds	R	R		R	R
<i>Rhynchosporium</i> Leaf Blotch		MC			C

### **Eyespot (*Oculimacula* spp.)**

Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

### **Septoria Leaf Spot and Glume Blotch (*Septoria tritici* and *Stagonospora nodorum*)**

Apply before disease is established in the crop. To protect the upper leaves and ear apply Cello at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

### **Powdery Mildew (*Blumeria graminis*)**

Apply Cello at the first signs of disease. Where disease pressure remains high application may be repeated.

### **Yellow Rust (*Puccinia striiformis*)**

Apply Cello at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

### **Brown Rust (*Puccinia recondita* and *P. hordei*)**

Apply Cello at the first signs of disease. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

### **Crown Rust (*Puccinia coronata*)**

Apply Cello at the first signs of disease. Cello controls Crown Rust in winter and spring oats. A second application may be made 2-3 weeks later if re-infection occurs. Applications made to established infections are likely to be less effective.

### **Tan Spot (*Pyrenophora tritici-repentis*)**

Apply Cello at the first signs of disease in spring or early summer. Where disease pressure remains high application may be repeated.

## Ear Disease Complex

Apply Cello soon after ear emergence until the end of flowering (GS 59-69) for control of *Fusarium* Ear Blight and reduction of Sooty Moulds. Control of ear diseases can result in cleaner, brighter ears.

Through the reduction of ear blight, Cello effectively reduces the level of the *Fusarium* mycotoxin deoxynivalenol (DON) in wheat grain. However, where *Fusarium* levels are high, the reduction achieved may not always be sufficient to ensure that DON levels fall below the statutory limit.

## Leaf Blotch (*Rhynchosporium commune*)

Apply Cello in spring at the first signs of disease. For severe infections a second application may be necessary 2-3 weeks later.

## Net Blotch (*Pyrenophora teres*)

Apply Cello at the first signs of disease in spring/early summer. For severe infections, a second application 2-3 weeks later will give most effective control when conditions remain favourable for disease development.

**CAUTION:** The possible development of disease strains resistant to Cello cannot be excluded or predicted. Where such resistant strains occur, Cello is unlikely to give satisfactory control.

## CROP SPECIFIC INFORMATION

Cello may be used on all commercial varieties of winter and spring barley, winter and spring oats, winter and spring wheat, winter rye and triticale

## Mixing

Thoroughly shake the pack before use.

Add the required quantity of Cello to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

## General

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.



## SAFETY PRECAUTIONS

### Operator Protection

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.

Wear suitable protective clothing (coveralls) when applying the product.

Wear suitable protective clothing (coveralls) and suitable protective gloves when in contact with contaminated surfaces.

When using do not eat, drink or smoke.

Wash any contamination from eyes immediately.

Wash hands and exposed skin before meals and after work.

If you feel unwell, seek medical advice (show the label where possible).

### Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m\* to surface water bodies.

\*To reduce this buffer zone please refer to PRCD Guidance – STRIPE (Surface water Tool for Reducing the Impact of Pesticides in the Environment)

Aim spray away from water

### Storage and Disposal

Keep in original container tightly closed in a safe place.

Keep out of reach of children.

Keep away from food, drink and animal feedingsuffs.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

PROTECT FROM FROST



cellosdsie

To access the **Safety Data Sheet** for this product scan the code or use the link below:

[www.bayercropscience.ie/sds/cello.pdf](http://www.bayercropscience.ie/sds/cello.pdf)

or alternatively contact your supplier

IE86259605a rA4

Bayer