



# Ironmax<sup>®</sup> PRO

## 15 kg

### Molluscicide

**For the control of slugs and snails in all edible and non-edible crops grown outdoors or under protection.**

Ready for use bait containing 24.2 g/kg ferric phosphate

For professional use only

Authorisation Holder:  
De Sangosse SA  
Bonnell - CS10005  
47480 Pont-du-Casse, France  
www.desangosse.fr

Marketing company:  
Bayer CropScience Ltd  
Bayer Ltd, The Atrium  
Blackthorn Rd  
Sandyford, Dublin 18

Freephone 1800 818534.  
For 24 hour emergency information contact Bayer CropScience Ltd.  
Telephone: 00800 1020 3333

#### SAFETY INFORMATION

#### IRONMAX<sup>®</sup> PRO

Contains Ferric phosphate (anhydrous) 24.2 g/kg.

Keep out of reach of children  
Do not eat, drink or smoke when using the product  
Wear protective gloves  
Collect spillage  
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for clean empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use

PCS No. 06513

888051 480a 1x1

# Ironmax<sup>®</sup> PRO

### SAFETY PRECAUTIONS

#### Operator protection

Wash hands after handling/using the product or working in a previously treated field.

#### Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

#### Storage and disposal

Store in a cool dry place.  
Keep away from food, drink and animal feedingstuffs.  
Do not use original packaging or empty container for any other purpose.  
IRONMAX PRO is a biological slug and snail control product for all edible and non-edible crops grown outdoors and under protection. IRONMAX PRO is a molluscicide that acts by ingestion. Once ingested, IRONMAX PRO blocks the pest's digestion processes.

### DIRECTIONS FOR USE

For use only as a molluscicide.

#### Warnings

Due to the mode of action of ferric phosphate, following consumption of the bait slugs usually retreat underground to die and there will be little evidence of dead slugs. The effects of treatment may therefore not be immediately visible. Effectiveness should therefore be measured by the decrease of feeding damage in the crop. The result of treatment with IRONMAX PRO can therefore only be assessed on the basis of reduced crop damage.

#### Crop specific information

| Crops                                       | Maximum individual dose | Maximum number of treatments | Pre-harvest interval |
|---|-------------------------|------------------------------|----------------------|
| All edible crops, outdoor and protected     | 7 kg/ha                 | 4 per crop                   | 3 days               |
| All non-edible crops, outdoor and protected | 7 kg/ha                 | 4 per crop                   | -                    |

#### Application

IRONMAX PRO may be applied by mechanical applicator; by hand or by hand-held equipment.  
Take care during application to prevent pellets resting on or lodging in leaves, florets or other parts of the plants.

#### Intervals and number of applications

Re-treatment may be required, especially when pellets have been consumed or when subsequent generations occur.  
Interval between applications: minimum 5 days.

#### Application timing

Best results are achieved when treatment takes place before damage occurs. For annual crops, it is advised to treat before emergence. Apply the product after seedbed preparation is complete.

#### Application timing should take into consideration:

- Pest detection or likelihood of activity
- Part of the crop targeted by the pest, whether it is the seeds, seedling plants or harvestable produce that are affected will define a specific strategy.

For further information on slug trapping and damage risk assessment, please refer to AHDB Information Sheet 04 'Integrated Slug Control' which can be found at: [www.ahdb.org.uk/slugcontrol](http://www.ahdb.org.uk/slugcontrol)

## 15 kg