



# Jettano®

3 L e

GROUP 7 FUNGICIDE

**A fungicide for the control  
of foliar diseases in wheat.**

An emulsifiable concentrate  
formulation containing  
100 g/L (10.3% w/w) bixafen  
and 100 g/L (10.3% w/w)  
fluopyram.



F

For Professional use only

#### Safety information

**JETTANO** UFI: WWG0-T0FG-900K-61VA

Contains 100 g/L (10.3% w/w) bixafen,  
100 g/L (10.3% w/w) fluopyram, Solvent Naphtha  
(petroleum), heavy aromatic, <1% naphthalene and  
N,N-Dimethyl decanamide.

#### **Danger**

Causes skin irritation.

May cause respiratory irritation.

May cause an allergic skin reaction.

May be fatal if swallowed and enters airways

Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Avoid breathing spray.

Wear protective gloves/protective clothing/eye  
protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

If Swallowed: Immediately call a POISON CENTER or doctor/  
physician.

If inhaled: Remove victim to fresh air and keep at rest in a position  
comfortable for breathing.

Dispose of contents/container to a licensed hazardous-waste  
disposal contractor or collection site except for triple rinsed empty  
containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply  
with the instructions for use.**

**PCS No. 07313**



IE91176046a rA2

## **SAFETY PRECAUTIONS**

### **Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

### **Environmental Protection**

Do not contaminate water with the product or its container. (Do not clean application

**Authorisation holder:** Bayer CropScience Limited  
230 Cambridge Science Park, Milton Road,  
Cambridge, CB4 0WB, United Kingdom

**Marketing Company:** Bayer CropScience Ltd,  
Bayer Ltd, 1st Floor, The Grange Offices, The  
Grange, Brewery Road, Stillorgan, Co. Dublin  
A94 H2K7 Freephone: 1800 818534

equipment near surface water. Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

### **Storage and Disposal**

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

Keep out of reach of children.

Keep away from food, drink and animal feedingstuffs.

PROTECT FROM FROST

### **For 24 hour emergency information contact**

**Bayer CropScience Ltd. Telephone:**

**+44 330 678 3382** (24 hr) (charged as a standard international call to the UK)

National Poisons Information Centre Dublin:

**01 809 2166**

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Jettano is a mixture of two SDHI fungicides from different chemical groups, a carboxamide and a benzamide, recommended for control of diseases on cereals.

### RATE OF USE

<b>Crop</b>	<b>Maximum individual/ total dose:</b>	<b>Maximum number of applications per crop</b>	<b>Latest time of application</b>
Winter and spring wheat, durum wheat spelt and triticale	1.25 litres product per hectare	One per crop	Before beginning of flowering, (GS 59)

### Other specific restrictions

The total number of foliar applications of products containing SDHI fungicides to any cereal crop must not exceed two. Mixtures of two or more SDHI fungicides do not provide an anti-resistance strategy. Each application of such a mixture counts as one SDHI application.

Apply SDHI fungicides always in mixtures.

The mixture partner:

- should provide satisfactory disease control when used alone on the target disease
- must have a different mode of action

Do not apply product prior to the beginning of stem elongation (BBCH 30)

**Method of application:** Tractor mounted/trailed sprayer.

A spray pressure of 2-3 bar is recommended. Apply Jettano in 150-400 litres per hectare water.

Apply as a medium spray quality.

## **CROPS**

Jettano may be used on all commercial varieties of winter and spring wheat.

## **RATE OF USE**

Apply Jettano at: 1.25 litre per hectare.

The maximum number of applications should be one per crop.

## **APPLICATION**

### **Water volume**

Apply Jettano in 150-400 litres per hectare water. The higher spray volumes are recommended where the crop is dense to ensure good penetration to the lower leaves and stem bases.

Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.

A spray pressure of 2-3 bar is recommended.

**Spray quality**

Apply as a MEDIUM spray quality (as defined by BCPC).

**Latest Permitted Timing**

Before beginning of flowering (BBCH 59).

**Mixing**

Thoroughly shake the pack before use.

Add the required quantity of Jettano to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

**General**

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

## DISEASES CONTROLLED

	Wheat	Barley
Septoria leaf blotch ( <i>Mycosphaerella graminicola</i> )	C	
Brown rust	C	
C = Control		

## APPLICATION TIMING

Only one application of Jettano is permitted, where disease pressure remains high follow up spray(s) with another effective product may be required.

### **Septoria Leaf Blotch (*Mycosphaerella graminicola*)**

Apply before disease is established in the crop up to ear emerged (GS 59). For severe infections a second application of another effective product may be necessary 2-3 weeks later.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective.

### **Brown Rust (*Puccinia triticina*, *P. recondita*)**

Apply Jettano at the first signs of disease. Applications made to established infections are likely to be less effective.

## **RESISTANCE STRATEGY**

Tank mixtures or alternation with fungicides having a different mode of action against the diseases present have been shown to protect against the development of resistant forms of disease.

Apply SDHI fungicides always in mixtures

The mixture partner:

- should provide satisfactory disease control when used alone on the target disease
- must have a different mode of action

No more than two applications of SDH inhibitors must be applied to the same crop.

Bixafen and fluopyram are SDH inhibitors.

Do not tank mix Jettano with other SDHI containing formulations.

Use Jettano as part of an Integrated Crop Management strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action and non-chemical methods of disease reduction.

**CAUTION:** The possible development of disease strains resistant to Jettano cannot be excluded or predicted. Where such resistant strains occur, Jettano is unlikely to give satisfactory control.

Jettano is a Registered Trade Mark of Bayer  
© Bayer CropScience Limited 2025



To access the **Safety Data Sheet** for this product  
scan the code or use the link below:

[www.bayercropscience.ie/sds/jettano.pdf](http://www.bayercropscience.ie/sds/jettano.pdf)

or alternatively contact your supplier